

HEART FAILURE WITH NORMAL AND REDUCED EJECTION FRACTION – ASSESSMENT AND SHARED CARE MANAGEMENT

Dr Rohit Khurana

Keywords: Heart Failure, shared care management, reduced ejection fraction, preserved ejection fraction, BNP, therapy

SFP2022; 48(1): X-X

INTRODUCTION

Heart failure (HF) is a common clinical syndrome resulting from any structural or functional cardiac disorder that impairs the ability of the ventricle to fill with or eject blood. HF may be caused by disease of the myocardium, pericardium, endocardium, heart valves, vessels, or by metabolic disorders.

HF due to left ventricular dysfunction is categorised into HF with reduced ejection fraction (with Left Ventricular Ejection Fraction (LVEF) ≤ 50 percent, known as HFrEF; also referred to as systolic HF) and HF with preserved ejection fraction (with LVEF > 50 percent; known as HFpEF; also referred to as diastolic HF¹).

A reduced LVEF in systolic heart failure is a powerful predictor of mortality. As many as 40-50 percent of patients with heart failure have diastolic heart failure with preserved left ventricular function.

Overall, there is no difference in survival between diastolic and systolic heart failure that cannot be attributed to ejection fraction. Patients with diastolic heart failure are more likely to be women, to be older, and to have hypertension, atrial fibrillation, and left ventricular hypertrophy, but no history of coronary artery disease.^{2,3} The pathogenesis of diastolic dysfunction involves abnormalities of active ventricular relaxation and passive ventricular compliance, which lead to ventricular stiffness and higher diastolic pressures. These pressures are transmitted through atrial and pulmonary venous systems, reducing lung compliance. A combination of decreased lung compliance and cardiac output leads to symptoms.²

DR ROHIT KHURANA

Senior Consultant, General and Interventional Cardiologist
The Harley Street Heart and Vascular Centre
Gleneagles Hospital

CLINICAL PRESENTATION AND DIAGNOSIS

Patients with HF usually present with the classic triad of symptoms – oedema, fatigue, and dyspnea. Other typical symptoms may include orthopnea, paroxysmal nocturnal dyspnea, reduced exercise tolerance, and increased time to recover from exercise. Less typical signs include nocturnal cough, wheezing, bloated feeling, loss of appetite, confusion (especially in the elderly), depression, palpitations, dizziness and syncope.⁴

However, symptoms are non-specific and non-sensitive, and therefore are less useful in discriminating HFrEF from HFpEF. In addition, atypical presentations should be considered when evaluating obese patients and older adults because of potentially different aetiology, clinical presentation, and outcome as compared to the general population.

Typical signs of HFrEF include elevated-jugular venous pressure, gallop rhythm, hepato-jugular reflux, and laterally displaced apical impulse. Less typical symptoms include weight gain, cachexia (tissue wasting), cardiac murmur, peripheral edema (ankle, sacral, scrotal), pulmonary rales, tachycardia, irregular pulse, tachypnea, hepatomegaly, ascites, cold extremities, oliguria, and narrow pulse pressure.⁴

The assessment of signs and symptoms is clinically significant to suggest the likelihood of HFrEF as well as to monitor response to therapy and stability overtime. In groups such as the obese, the elderly and those with chronic lung disease, being confident of the symptoms and signs may be more difficult.

Persistent symptoms despite treatment often suggest the need for additional therapy and worsening of symptoms often suggest serious development and the need for prompt medical attention.

Heart failure is much more likely in the presence of a relevant medical history suggesting an increased risk of cardiac damage. In advanced economies such as Singapore, nearly 60 percent of patients diagnosed with acute heart failure have underlying coronary artery disease and in patients with acute coronary syndromes, myocardial ischaemia is often a precipitant risk factor, especially for de novo heart failure.⁵ A prior diagnosis of heart failure, diabetes, hypertension, valvular heart disease, advanced age, the male sex, and obesity have all been found to predict fluid overload typical of congestive heart failure.

The symptoms and signs of heart failure are the consequence of systemic and pulmonary congestion, which result from increased left ventricular filling pressures. Even in the absence of overt clinical congestion, hemodynamic congestion may still occur, and this also predicts a worse mortality and re-

hospitalisation rate. Heart failure symptoms can occur with preserved or reduced ejection fraction (systolic or diastolic heart failure). The New York Heart Association classification system is the simplest and most widely used method to gauge symptom severity (Table 1). The classification system is a well-established predictor of mortality and can be used at diagnosis and to monitor treatment response.

Table 1: American College of Cardiology/American Heart Association Recommendations by Stage of Heart Failure¹³

STAGE	RECOMMENDATION
A: Heart failure risk factors	Guideline-directed treatment of hypertension and hyperlipidemia
B: Diastolic dysfunction without symptoms	Treat hypertension with thiazide diuretics, ACE inhibitors, or nondihydropyridine calcium channel blockers
C: Symptomatic heart failure with preserved ejection fraction and hypertension	Treat volume overload with diuretics; consider use of beta blockers, ACE inhibitors
C: Symptomatic heart failure with preserved ejection fraction without hypertension	Treat volume overload with diuretics

ACE = angiotensin-converting enzyme; ARB = angiotensin receptor blocker.

Information from references 3 and 18.

ASSESSMENT

In patients with non-acute onset, presenting with symptoms and signs of heart failure in the primary care setting, the probability of heart failure should be determined by the history and symptoms supported by findings on clinical examination. Blood tests, an ECG and a chest x-ray should be requested on all patients. During these steps, at least one element should be positive to consider the diagnosis and subsequently plasma natriuretic peptides and an echocardiogram should be requested. The diagnostic assessment, in accordance with the ESC 2016 guidelines, is summarised in Table 2.⁴

Table 2: Diagnostic assessment in patient with suspected heart failure (non-acute onset) – according to ESC guidelines⁴

Diagnostic assessment in patient with acute heart failure adapted from the ESC guidelines (2016)¹

1. Measurement of plasma natriuretic peptide level (BNP, NT-proBNP or MR-proANP)*

2.

- a) 12-lead ECG;
- b) chest X-ray†
- c) laboratory assessments: cardiac troponins, BUN (or urea), creatinine, electrolytes (sodium, potassium), glucose, complete blood count, liver function tests and TSH.

3. Echocardiography‡

* to help in the differentiation of acute heart failure from non-cardiac causes of acute dyspnea

† to assess signs of pulmonary congestion and detect other cardiac or non-cardiac diseases that may cause or contribute to the patient's symptoms

‡ recommended immediately in haemodynamically unstable patients with acute heart failure and within 48 hours when cardiac structure and function are either not known or may have changed since previous studies.

Brain Natriuretic Peptide (BNP)

BNP and N-terminal pro-BNP (the cleaved inactive N-terminal fragment of the BNP precursor) levels can be used to evaluate patients with shortness of breath suspected of having heart failure.⁵ BNP is secreted by the atria and ventricles in response to stretching or increased wall tension. The hormone then causes fluid and sodium loss in the urine and mild vasodilation. BNP levels increase with age, is higher in women and patients of African origin, and can be elevated in patients with renal failure. BNP appears to have better reliability than N-terminal pro-BNP,⁶ especially in older populations. Guidelines from the American College of Cardiology/American Heart Association (ACC/AHA) and European Society of Cardiology (ESC) recommend the use of natriuretic peptides for assessment of patients with symptoms of heart failure.

Most dyspneic patients with HF have plasma BNP values >400 pg/mL, while values <100 pg/mL have a very high negative predictive value for HF as a cause of dyspnea.⁷ In the range between 100–400 pg/mL, plasma BNP concentrations are not very sensitive or specific for detecting or excluding HF. As BNP levels increase, the specificity increases and likewise the likelihood of a heart failure diagnosis. BNP levels also increase in level according to New York Heart Association classification. Elevations in plasma BNP can establish the presence of HF due to diastolic dysfunction with similar accuracy to systolic dysfunction. However, the values do not differentiate between systolic and diastolic dysfunction.

BNP levels are strong predictors of mortality at two to three months and cardiovascular events in acute heart failure, specifically when BNP level is >200 pg/mL or N-terminal pro-BNP level is >5,180 pg/mL.⁷ Limited evidence supports monitoring reduction of BNP levels in the acute and outpatient settings. A 30 to 50 percent reduction in BNP level at hospital discharge showed improved survival and reduced re-hospitalisation rates. Optimising management for outpatient targets of a BNP level <100 pg/mL and an N-terminal pro-BNP level <1,700 pg/mL showed improvement in decompensations, hospitalisations, and mortality events.⁷

Echocardiography

Echocardiography should be performed in all patients with new onset HF and can provide important information about ventricular size and function.⁸ The sensitivity and specificity of two-dimensional echocardiography for the diagnosis of systolic dysfunction are as high as 80 and 100 percent, respectively. Valvular structure and function in valve disease can be characterised.

A number of other important findings can be detected. Regional wall motion abnormalities are compatible with coronary artery disease. Pericardial thickening may be indicative of constrictive pericarditis. Infiltrative

cardiomyopathies are associated with an abnormal myocardial texture. Left ventricular diastolic function can be assessed by estimation of the pulmonary capillary wedge pressure via the ratio of tissue Doppler of early mitral inflow velocity (E) to early diastolic velocity of the mitral annulus (e'). An E/e' ratio of >15 suggests a Pulmonary capillary wedge pressure (PCWP) of >15 mmHg when e' is the mean of medial and lateral mitral annulus early diastolic velocities. There are limitations to the use of the E/e' ratio, which are beyond the scope of this article. Right atrial and pulmonary artery pressures are determined by the peak velocity of tricuspid regurgitation on Doppler echocardiography.

These findings correlate with the pulmonary artery wedge pressure, regardless of the etiology of HF or severity of tricuspid regurgitation; they can be used to assess changes in left ventricular filling pressures resulting from therapy.

Treadmill Exercise Testing

Exercise testing should be part of the initial evaluation of virtually all patients with HF. In addition to detection of ischemic heart disease, assessment of exercise capacity can be used for risk stratification and determining prognosis. Serial measurements can also assess the efficacy of therapy and clinical stability of patients over time.

With severe HF, measurement of the maximal oxygen uptake (VO_{2max}) provides an objective estimate of the functional severity of the myocardial dysfunction.⁹ VO_{2max} is one of the best indices of prognosis in patients with symptomatic HF. However, peak VO_{2max} and exercise capacity can be affected by factors other than cardiac status, including deconditioning, pulmonary disease, and anaemia. One advantage of measuring VO_{2max} directly is that cardiac and non-cardiac causes of impaired exercise can be distinguished by assessing the anaerobic threshold and related indices.

PRINCIPLES OF MANAGEMENT

The 2016 ESC guidelines recommend both lifestyle interventions and pharmacological therapies.⁴ HF with reduced ejection fraction has well-validated therapies to reduce morbidity (i.e., reducing symptoms, improving health-related quality of life and functional status, decreasing the rate of hospitalisation), and to reduce mortality. In contrast, HF with preserved ejection fraction lacks evidence-based treatment recommendations.

Diuretics are essential in relieving symptoms, but it has yet to be definitively established whether they provide long-term prognostic benefit.¹⁰ They provide immediate relief of symptoms and help manage the chronic fluid status in stable patients, irrespective of whether it is HFrEF or HFpEF.


It is imperative that patients with heart failure understand their condition and are involved in their management decisions. Lifestyle interventions can improve patients' quality of life and prevent exacerbations. The role of dietary

salt restriction and the importance of regular exercise to increase functional capacity should be reinforced. For patients with more advanced heart failure, daily weight measurements and fluid restriction with close supervision may be necessary. Formalised cardiac rehabilitation that combines exercise with ongoing educational and psychological support has proven benefits.

HF with reduced ejection fraction

Management of heart failure with reduced ejection fraction (HFrEF) includes management of contributing factors and associated conditions, lifestyle modification, pharmacologic therapy, and, if indicated, device therapy (Figure 1).

Figure 1. Therapeutic algorithm for a patient with symptomatic heart failure with reduced ejection fraction⁴



Cardiac resynchronisation therapy is indicated with patients with an ejection fraction of <35 percent and QRS duration of >130 ms. Contributing factors comprise hypertension, myocardial ischemia or infarction, diabetes mellitus, thyroid dysfunction, and infection. Coronary disease and diabetes mellitus have become increasingly responsible for HF while hypertension and valve disease have become less common due to improvements in detection and therapy. However, almost all patients with HFrEF will require diuretics on a chronic basis to alleviate symptoms and signs of congestion.¹¹

Other treatment generally starts with ACE inhibitors or angiotensin receptor blockers (ARBs), followed by beta-blockers. Beta blockers can also provide relief in patients with ischemic heart disease presenting with angina and rate control in those with atrial fibrillation. If symptoms persist, as based on the NYHA functional classification of

II-IV, mineralocorticoid receptor antagonists (MRA such as spironolactone or eplerenone may be added. Ivabradine may also be an option for some patients with HFrEF, if they are in sinus rhythm and their resting heart rate is >70 bpm. *Improved mortality* has been demonstrated with beta blockers, ACE inhibitors, ARNI, hydralazine plus nitrate, and MRAs, with limited evidence of survival benefit for diuretics.

NOVEL TREATMENT OPTIONS IN HFREF

One new treatment strategy is the angiotensin receptor neprilysin inhibitor (ARNI), which comprises valsartan and sacubitril. This drug serves to reduce sympathetic tone, aldosterone levels, and sodium retention through inhibition of the overactive renin angiotensin system while simultaneously potentiating protective vasoreactive neuropeptides.

The first (and currently only commercially available) ARNI, formerly known as LCZ696 and marketed as Entresto, was evaluated in the PARADIGM-HF trial in comparison to enalapril 10mg bd. After 27 months of follow-up, the trial was stopped early due to positive interim results.¹² All-cause mortality was 17 percent with the ARNI, as compared to 19.8 percent with enalapril, in HFrEF patients maintained on optimal background HF therapy. This yielded a hazard ratio of 0.84 ($p < 0.001$) and a number needed to treat of 32.

The position of ARNI prescribing in primary care has also not been firmly established. Cardioselective beta-blockers may be prescribed in patients with co-morbid chronic obstructive pulmonary disease (COPD). More recently, the EMPA-REG OUTCOME and the DECLARE TIMI-58 trial reported that empagliflozin and dapagliflozin, respectively, which are both inhibitors of the sodium glucose co-transporter (SGLT-2) in the kidney when added to metformin, had beneficial prognostic cardiovascular outcomes compared to placebo in patients with cardiovascular risk factors and/or disease.¹² Subgroup analysis suggested the benefits were consistent for patients with and without HF.

HF with preserved ejection fraction

In contrast with treatment of heart failure with HFrEF, there are fewer randomised controlled trials (RCTs) of patients with HFpEF. The ACC/AHA recommends using a stage-based approach to guide treatment of HFpEF.¹³ The ESC recommends diuretics for treating fluid overload in patients with HFpEF.⁴ However, it makes no recommendation regarding other medications for treatment. It recommends identifying and treating cardiovascular and non-cardiovascular comorbidities, because most deaths and hospitalisations in patients with HFpEF are not due to chronic heart failure. The ACC/AHA and ESC recommend combined endurance and resistance training for patients with HFpEF to improve exercise capacity, physical functioning, and diastolic function.

Hypertension should be treated according to appropriate guidelines. Although RCTs of several medications showed fewer heart failure hospitalisations, this benefit was offset by increases in hospitalisations for other reasons. Thus, in the absence of hypertension, the evidence does not support treating patients with HFpEF with any medication except diuretics. Additionally, RCTs of angiotensin receptor blockers, nitrates, and spironolactone raise concerns about adverse effects, and physicians should avoid using these medications if possible. Similarly, the use of digoxin should be avoided in patients 65 years and older. Comorbid atrial fibrillation or coronary artery disease should be treated.

Data from the PARAMOUNT trial comparing valsartan-sacubitril with valsartan in HFpEF showed that ARNI reduced NT-proBNP levels, left atrial volume index, and increased eGFR more so than valsartan alone and independent of its systolic blood pressure lowering effect. The long-term clinical outcomes of ARNI in HFpEF are being further investigated in the ongoing PARAGON trial.

SHARED CARE MANAGEMENT

Often, patients with HFrEF are managed in the hospital outpatient or specialist clinics for three to six months after diagnosis, to titrate medication to optimal doses. The optimal duration of these hospital-based programmes before transitioning patient care to the community has not been established, nor whether all or only higher risk patients benefit. However, there is evidence that, given the complexity of the HF syndrome and its co-morbidities, close collaboration between hospital and primary care practitioners is crucial in order to provide optimal, integrated care and improved outcomes.¹⁴ It is important to realise that there is a wide variation in the disease trajectory of HF and the majority of patients do not generally follow a gradual decline. Some feel and function well but succumb to sudden fatality, while other patients may display improved functioning after a period of poor quality of life. Diverse and multiple co-morbidities further complicate the disease trajectory, warranting regular monitoring.

REFERENCES

- McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2012 Jul;33(14):1787-847. doi: 10.1093/eurheartj/ehs104. Epub 2012 May 19. Erratum in: *Eur Heart J*. 2013 Jan;34(2):158. PMID: 22611136.
- Sharma K, Kass DA. Heart failure with preserved ejection fraction: mechanisms, clinical features, and therapies. *Circ Res*. 2014 Jun 20;115(1):79-96. doi: 10.1161/CIRCRESAHA.115.302922. PMID: 24951759; PMCID: PMC4146618.
- Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of

- the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail.* 2016 Aug;18(8):891-975. doi: 10.1002/ehf.592. Epub 2016 May 20. PMID: 27207191.
4. Chen WC, Tran KD, Maisel AS. Biomarkers in heart failure. *Heart.* 2010 Feb;96(4):314-20. doi: 10.1136/hrt.2008.151639. PMID: 20194212.
 5. Kelder JC, Cramer MJ, van Wijngaarden J, van Tooren R, Mosterd A, Moons KG, et al. The diagnostic value of physical examination and additional testing in primary care patients with suspected heart failure. *Circulation.* 2011 Dec 20;124(25):2865-73. doi: 10.1161/CIRCULATIONAHA.111.019216. Epub 2011 Nov 21. PMID: 22104551.
 6. Maisel A, Mueller C, Adams Jr K, Anker SD, Aspromonte N, Cleland JG, et al. State of the art: using natriuretic peptide levels in clinical practice. *Eur J Heart Fail.* 2008 Sep;10(9):824-39. doi: 10.1016/j.ejheart.2008.07.014. Epub 2008 Aug 29. PMID: 18760965.
 7. Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, et al. 2009 focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation. *Circulation.* 2009 Apr 14;119(14):e391-479. doi: 10.1161/CIRCULATIONAHA.109.192065. Epub 2009 Mar 26. Erratum in: *Circulation.* 2010 Mar 30;121(12):e258. PMID: 19324966.
 8. Ingle L. Prognostic value and diagnostic potential of cardiopulmonary exercise testing in patients with chronic heart failure. *Eur J Heart Fail.* 2008 Feb;10(2):112-8. doi: 10.1016/j.ejheart.2007.12.011. Epub 2008 Feb 5. PMID: 18255336.
 9. von Lueder TG, Atar D, Krum H. Diuretic use in heart failure and outcomes. *Clin Pharmacol Ther.* 2013 Oct;94(4):490-8. doi: 10.1038/clpt.2013.140. Epub 2013 Jul 12. PMID: 23852396.
 10. McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med.* 2014 Sep 11;371(11):993-1004. doi: 10.1056/NEJMoa1409077. Epub 2014 Aug 30. PMID: 25176015.
 11. Giugliano D, Meier JJ, Esposito K. Heart failure and type 2 diabetes: From cardiovascular outcome trials, with hope. *Diabetes Obes Metab.* 2019 May;21(5):1081-1087. doi: 10.1111/dom.13629. Epub 2019 Feb 14. PMID: 30609236.
 12. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, et al. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation.* 2013 Oct 15;128(16):1810-52. doi: 10.1161/CIR.0b013e31829e8807. Epub 2013 Jun 5. PMID: 23741057.
 13. Luttik ML, Jaarsma T, van Geel PP, Brons M, Hillege HL, Hoes AW, et al. Long-term follow-up in optimally treated and stable heart failure patients: primary care vs. heart failure clinic. Results of the COACH-2 study. *Eur J Heart Fail.* 2014 Nov;16(11):1241-8. doi: 10.1002/ehf.173. Epub 2014 Oct 10. PMID: 25302753.

LEARNING POINTS

- **It is imperative that patients with heart failure understand their condition and are involved in their management decisions; lifestyle interventions can improve patients' quality of life and prevent exacerbations.**
- **Given the complexity of the HF syndrome and its co-morbidities, close collaboration between hospital and primary care practitioners is crucial in order to provide optimal, integrated care and improved outcomes.**
- **Recent trials (EMPA-REG OUTCOME and the DECLARE TIMI-58) suggest that both empagliflozin and dapagliflozin, which are inhibitors of the sodium glucose co-transporter (SGLT-2) in the kidney, had beneficial prognostic cardiovascular outcomes when added to metformin compared to placebo in patients with cardiovascular risk factors and/or disease.**
- **Heart failure is increasingly common. Timely and accurate diagnosis is important since treatments can improve symptoms and improve prognosis.**
- **There are parallels with the diagnostic evaluation of heart failure with both preserved and reduced ejection fraction. Management approached overlap but the syndromes also require distinct treatment strategies.**
- **Primary care has a crucial responsibility to oversee the health status of a heart failure patient and associated co-morbid conditions using evidence-based pharmacotherapy and lifestyle interventions.**