

DIETARY INTERVENTIONS FOR WEIGHT LOSS

Ms Izabela Kerner

ABSTRACT

Obesity is a complex and chronic condition that requires continuing care. A variety of diet plans are available for use in the clinical setting. Exactly what type of diet is the most beneficial remains up for debate. Numerous clinical trials have been carried out over the years to compare an array of dietary interventions for weight loss, including calorie-restricted diets, altered macronutrient composition diets, or specific dietary patterns. This paper will provide an overview of some of the evidence-based dietary interventions for clinical practice.

Keywords: Obesity, weight loss, dietary interventions, low-carbohydrate diet, energy restriction

SFP2025; 51(5): 20-23

INTRODUCTION

Obesity is a serious global epidemic associated with numerous metabolic complications, including type 2 diabetes (T2DM), hypertension, cardiovascular disorders, and several cancers. The aetiology of obesity is multifactorial, involving an interplay of genetic, biological, environmental, social, cultural, and behavioural factors. Even though a successful weight loss strategy should be individualised and address all the underlying causes of obesity, dietary and lifestyle interventions remain the cornerstone of treatment. However, the optimal dietary approach to weight loss is still a subject of debate amongst experts, healthcare professionals, and the public, as studies have failed to demonstrate the superiority of one diet plan over another in the long term. Comprehensive assessment of an individual's dietary habits and lifestyle should be the first step in deciding on the best dietary intervention for weight loss and avoiding a one-size-fits-all approach. There is evidence that even modest weight loss of 5–8 percent body weight achieved with diet and lifestyle interventions improves glycaemia, blood pressure, lipid profile, mobility, and quality of life.^{1,2} Greater weight loss might be needed to produce health benefits in individuals with morbid obesity or multiple comorbidities.

MS IZABELA KERNER
 Consultant Dietitian, APD, ADS
 Digestive Centre, Mount Elizabeth Hospital
 Member, Singapore Nutrition and Dietetics Association (SNDA)
 Past President of SNDA (April 2018-March 2022)

ENERGY RESTRICTION DIET INTERVENTIONS

Energy-restricted diets range from continuous or intermittent diet plans with moderate to severe caloric deficit depending on the health status and weight loss goals of a patient or client. A calorie deficit of at least 500–750 per day is recommended to achieve a weight loss of 0.5–1.0 kg per week, which is the standard approach in clinical practice. These continuous energy restriction diet plans are formulated using healthy eating and exercise principles incorporating individual food preferences and behaviour modification to increase compliance. However, specially formulated meals or partial liquid meal replacements have also been employed to help achieve the necessary caloric deficit to drive weight loss in certain individuals after considering their health status, dietary preferences, and cost.

On the other hand, Low and Very Low Energy Diets (VLED) based on total meal replacements restrict energy intake to about 800 calories per day or less, which results in a more rapid weight loss and improvement of obesity related comorbidities including remission of type 2 diabetes.³⁻⁵ The use of total meal replacements helps to deliver adequate levels of essential nutrients and improves adherence. VLEDs have been underutilised in clinical practice in view of concerns over the potential loss of lean body mass, risk of precipitating eating disorder, overall safety, and subsequent weight regain due to rapid weight loss. However, when clinically supervised, there is no evidence of VLED causing eating disorders or resulting in worse outcomes on knee strength, handgrip strength, or bone density compared to moderate energy-restricted diets in short-term studies lasting 3–6 months.⁶⁻⁸

The use of VLEDs in supervised conditions for up to three months in patients who fail to meet weight or metabolic targets with a standard approach is gaining support from institutions such as the National Institute for Health and Clinical Excellence and clinicians. A structured, individualised weight maintenance programme with gradual readjustment to normal eating after VLED is necessary to maintain weight loss and metabolic benefits.^{5,9,10} Further research is needed to assess the long-term impact of VLED plans on body composition, weight maintenance, and metabolic outcomes.

Intermittent energy restriction (IER) is another dietary approach for weight loss that involves periods of fasting and eating or Intermittent Fasting (IF) with or without calorie restriction. Popular types of IF include alternate-day fasting (ADF), the 5:2 fast (five days of normal eating and two days of restricted eating per week), and time-restricted eating (TRE).¹¹ In the TRE trials, the fasting window varies from 12 to 20 hours with an *ad libitum* diet during the eating period.

The key appeal of IF is high compliance, easy sustainability compared to continuous daily caloric restriction, and other potential health benefits such as improvements in cognitive function, metabolic health, and longevity demonstrated by animal studies.¹² However, feelings of hunger may be more pronounced during IER.

The evidence from human trials suggests that IF is safe for most healthy adults and achieves comparable weight loss and metabolic improvements to continuous energy restriction.^{13,14} In recent meta-analyses, both intermittent and continuous energy restriction resulted in a similar weight loss, weight maintenance, and improvements in cardiovascular risk factors.^{15,16} However, IF may be more effective as a weight loss intervention for people with a higher BMI.¹⁷ Many studies on IF in humans are short-term and involve a small number of subjects. Furthermore, with the focus of IER on the timing of eating rather than on food choices, some individuals might not achieve a desirable diet quality. It is therefore important to counsel patients about making healthy food choices when they are not fasting to improve their health further or prevent potential nutrient deficiencies. Longer-term trials are needed to further address the safety and efficacy of IF for different patient groups to help build confidence in recommending these IER plans.

ALTERED MACRONUTRIENT COMPOSITION DIETS

Low and very low carbohydrate diets, often referred to as “keto” diets, have gained popularity amongst healthcare professionals and the public as an effective tool for weight loss and a means to reduce metabolic complications associated with overweight and obesity. However, the terms “low carbohydrate diet”, “very low carbohydrate diet”, and “ketogenic diet” are often used interchangeably with lack of consensus in the literature on their definition. Low carbohydrate diets are defined as providing less than 26 percent of calories from carbohydrates (50–130 g per day) while very low carbohydrate diets limit carbohydrate calories to less than 10 percent (20–50 g per day).¹⁸

The ketogenic diet was initially developed to treat severe epilepsy in infants and children under medical supervision. It is a very low carbohydrate high-fat diet plan, resulting in a state of ketosis where fat instead of glucose is being burnt for fuel. True ketogenic diets used in clinical settings can limit carbohydrates to as little as 5 percent of calories, primarily from non-starchy vegetables, and provide up to 85 percent calories from fat with enough protein to preserve lean body mass but maintain ketosis.

However, when used as a tool for weight loss, these “keto” diet plans vary in the proportion of fat, protein, and carbohydrates they provide. The standard ketogenic diet usually provides 60–70 percent calories from fat, 20–30 percent from protein and up to 10 percent from carbohydrates. However, there is an individual variation in the level of carbohydrate and protein intake that is compatible with ketosis, thus diet personalisation is required. “Keto flu” is a frequent side

effect of a keto diet, which can include light-headedness, fatigue, headaches, nausea, and constipation, in particular during the adaptation phase. Multivitamin, mineral, and fibre supplements can be considered in some individuals to reduce side-effects. The nutritional adequacy of “keto” diet plans will depend on the overall diet composition and the nutrient sources.

A recent review of evidence on low and very low carbohydrate diets found them to be effective but not superior to other weight-loss diets.¹⁹ There was no difference in weight loss between lower carbohydrate (4–45 percent calories)/higher fat (30–75 percent calories) diets compared to higher carbohydrate (50–65 percent calories)/lower fat (20–25 percent calories) diets when protein and energy levels were kept the same.^{18,20} However, studies in patients who were overweight and diabetic following a low carbohydrate diet showed improvements in triglycerides and HDL cholesterol levels, insulin sensitivity, and glycaemic control with mixed effects on LDL cholesterol.^{21,22}

One advantage that “keto diets” may offer is controlling the cravings and hunger often reported with other diet plans. A review published in 2015 found that individuals adhering to a ketogenic diet reported significantly less hunger and desire to eat compared with baseline.²³ Even though well-formulated ketogenic diets may offer short-term health benefits in some individuals, they are difficult to sustain, and long-term risks and benefits are not fully understood in the absence of long-term studies.

SPECIFIC DIETARY PATTERNS

Diets focusing on dietary patterns such as the Dietary Approaches to Stop Hypertension (DASH) diet, Mediterranean diets, and Plant-Based Diets (PBD) have also been studied in weight loss trials.

The DASH diet recommends specific servings of different food groups depending on daily caloric needs. It focuses on whole grains, fruits and vegetables, fat-free and low-fat dairy, lean meat, fish, and poultry. A meta-analysis revealed that overweight and obese adults on the DASH diet lose more weight than controls following a standard diet in studies ranging from 8–24 weeks.²⁴ Calorie-restricted DASH diets led to even greater weight loss when compared to other low energy diets.

Mediterranean diets emphasise the intake of vegetables, fruit, legumes, nuts, whole grains, and olive oil as the main source of fat, with moderate amounts of fish and poultry, low intake of red meat, and moderate consumption of wine. Meta-analysis of RCTs found that energy-restricted Mediterranean diets achieve as much or more weight loss than low-carbohydrate and low-fat diets with or without energy restriction among overweight and obese adults when followed for at least six months.²⁵

A variety of PBDs including vegan and vegetarian diets have been investigated for their beneficial effects on weight loss

and associated comorbidities. In a recent systematic review, PBDs were found to be effective in reducing weight and waist circumference in individuals with T2DM, especially in studies with a duration of at least 16 weeks.²⁶ PBDs are characterised by high intake of dietary fibre from wholegrains, vegetables, and legumes, and low glycaemic index, which may enhance satiety and improve glycaemic control.

The DASH, Mediterranean diet, and PBDs can be safe, effective, and sustainable weight loss eating plans that also improve metabolic complications.

CONCLUSION

A variety of dietary approaches, with sufficient reduction in energy intake and high level of dietary adherence, can produce weight loss. IER and ketogenic diets are gaining popularity and have shown superiority in reducing body fat and improving obesity-related metabolic complications in the short term. However, studies have failed to establish the superiority of one diet plan over another in the long term. Weight loss interventions must address the underlying causes of overweight and obesity and facilitate sustainable behavioural and lifestyle changes to prevent and manage relapse.

REFERENCES

- Ryan DH, Yockey SR. Weight Loss and Improvement in Comorbidity: Differences at 5%, 10%, 15%, and Over. *Curr Obes Rep.* 2017 Jun;6(2):187-194. doi: 10.1007/s13679-017-0262-y. PMID: 28455679; PMCID: PMC5497590.
- Hamman RF, Wing RR, Edelstein SL, et al. Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes Care.* 2006 Sep;29(9):2102-7. doi: 10.2337/dc06-0560. PMID: 16936160; PMCID: PMC1762038.
- Lean ME, Leslie WS, Barnes AC, et al. Primary care-led weight management for remission of type 2 diabetes (DiRECT): an open-label, cluster-randomised trial. *Lancet.* 2018 Feb 10;391(10120):541-551. doi: 10.1016/S0140-6736(17)33102-1. Epub 2017 Dec 5. PMID: 29221645.
- Satter N, Welsh P, Leslie WS, et al. Dietary weight-management for type 2 diabetes remissions in South Asians: The South Asian diabetes remission randomised trial for proof-of-concept and feasibility (STANDbY). *Lancet.* 2022 Nov;9:1-15.
- Churuangsu C, Hall J, Reynolds A, Griffin SJ, Combet E, Lean ME. Diets for weight management in adults with type 2 diabetes: an umbrella review of published meta-analyses and systematic review of trials of diets for diabetes remission. *Diabetologia.* 2022 Jan;65(1):14-36. Epub 2021 Nov 17. PMID: 34796367; PMCID: PMC8660762.
- da Luz FQ, Hay P, Gibson AA, et al. Does severe dietary energy restriction increase binge eating in overweight or obese individuals? A systematic review. *Obes Rev.* 2015 Aug;16(8):652-65. doi: 10.1111/obr.12295. Epub 2015 Jun 11. PMID: 26094791.
- Zibellini J, Seimon RV, Lee CM, Gibson AA, Hsu MS, Sainsbury A. Effect of diet-induced weight loss on muscle strength in adults with overweight or obesity - a systematic review and meta-analysis of clinical trials. *Obes Rev.* 2016 Aug;17(8):647-63. doi: 10.1111/obr.12422. Epub 2016 Apr 29. PMID: 27126087.
- Zibellini J, Seimon RV, Lee CM, et al. Does Diet-Induced Weight Loss Lead to Bone Loss in Overweight or Obese Adults? A Systematic Review and Meta-Analysis of Clinical Trials. *J Bone Miner Res.* 2015 Dec;30(12):2168-78. doi: 10.1002/jbmr.2564. Epub 2015 Jun 16. PMID: 26012544.
- Lean M, Hankey C. Keeping it off: the challenge of weight-loss maintenance. *Lancet Diabetes Endocrinol.* 2018 Sep;6(9):681-683. doi: 10.1016/S2213-8587(17)30405-9. Epub 2018 Jan 19. PMID: 29371077.
- McCombie L, Brosnahan N, Ross H, et al. Filling the intervention gap: service evaluation of an intensive nonsurgical weight management programme for severe and complex obesity. *J Hum Nutr Diet.* 2019 Jun;32(3):329-337. doi: 10.1111/jhn.12611. Epub 2018 Nov 22. PMID: 30565772.
- Stockman MC, Thomas D, Burke J, Apovian CM. Intermittent Fasting: Is the Wait Worth the Weight? *Curr Obes Rep.* 2018 Jun;7(2):172-185. doi: 10.1007/s13679-018-0308-9. PMID: 29700718; PMCID: PMC5959807.
- Longo VD, Mattson MP. Fasting: molecular mechanisms and clinical applications. *Cell Metab.* 2014 Feb 4;19(2):181-92. doi: 10.1016/j.cmet.2013.12.008.
- Rynders CA, Thomas EA, Zaman A, Pan Z, Catenacci VA, Melanson EL. Effectiveness of Intermittent Fasting and Time-Restricted Feeding Compared to Continuous Energy Restriction for Weight Loss. *Nutrients.* 2019 Oct 14;11(10):2442. doi: 10.3390/nu11102442. PMID: 31614992; PMCID: PMC6836017.
- Wilkinson MJ, Manoogian ENC, Zadorian A, et al. Ten-Hour Time-Restricted Eating Reduces Weight, Blood Pressure, and Atherogenic Lipids in Patients with Metabolic Syndrome. *Cell Metab.* 2020 Jan 7;31(1):92-104.e5. doi: 10.1016/j.cmet.2019.11.004. Epub 2019 Dec 5. PMID: 31813824; PMCID: PMC6953486.
- Allaf M, Elghazaly H, Mohamed OG, et al. Intermittent fasting for the prevention of cardiovascular disease. *Cochrane Database Syst Rev.* 2021 Jan 29;1:CD013496. doi: 10.1002/14651858.CD013496.pub2. PMID: 33512717.
- Cioffi I, Evangelista A, Ponzio V, et al. Intermittent versus continuous energy restriction on weight loss and cardiometabolic outcomes: a systematic review and meta-analysis of randomised controlled trials. *J Transl Med.* 2018 Dec 24;16(1):371. doi: 10.1186/s12967-018-1748-4. PMID: 30583725; PMCID: PMC6304782.
- Yang F, Liu C, Liu X, et al. Effect of Epidemic Intermittent Fasting on Cardiometabolic Risk Factors: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Front Nutr.* 2021 Oct 18;8:669325. doi: 10.3389/fnut.2021.669325.
- Practice-based evidence in Nutrition (PEN) Global Resource for Nutrition Practice. Dietitians of Canada 2005-2021.
- Kirkpatrick CF, Bolick JP, Kris-Etherton PM, et al. Review of current evidence and clinical recommendations on the effects of low-carbohydrate and very-low-carbohydrate (including ketogenic) diets for the management of body weight and other cardiometabolic risk factors: A scientific statement from the National Lipid Association Nutrition and Lifestyle Task Force. *J Clin Lipidol.* 2019 Sep-Oct;13(5):689-711.e1. doi: 10.1016/j.jacl.2019.08.003. Epub 2019 Sep 13. PMID: 31611148.
- Kosinski C, Jornayvaz FR. Effects of Ketogenic Diets on Cardiovascular Risk Factors: Evidence from Animal and Human Studies. *Nutrients.* 2017 May 19;9(5):517. doi: 10.3390/nu9050517. PMID: 28534852; PMCID: PMC5452247.
- Abbasi J. Interest in the Ketogenic Diet Grows for Weight Loss and Type 2 Diabetes. *JAMA.* 2018 Jan 16;319(3):215-217. doi: 10.1001/jama.2017.20639. PMID: 29340675.
- Saslow LR, Daubenmier JJ, Moskowitz JT, et al. Twelve-month outcomes of a randomised trial of a moderate-carbohydrate versus very low-carbohydrate diet in overweight adults with type 2 diabetes mellitus or prediabetes. *Nutr Diabetes.* 2017 Dec 21;7(12):304. doi: 10.1038/s41387-017-0006-9. PMID: 29269731; PMCID: PMC5865541.
- Gibson AA, Seimon RV, Lee CM, et al. Do ketogenic diets really suppress appetite? A systematic review and meta-analysis. *Obes Rev.* 2015 Jan;16(1):64-76. doi: 10.1111/obr.12230. Epub 2014 Nov 17. PMID: 25402637.
- Soltani S, Shirani F, Chitsazi MJ, Salehi-Abargouei A. The effect of dietary approaches to stop hypertension (DASH) diet on weight and body composition in adults: a systematic review and meta-analysis of randomised controlled clinical trials. *Obes Rev.* 2016;17:442-454. doi: 10.1111/obr.12391

25. Esposito K, Kastorini CM, Panagiotakos DB, Giugliano D. Mediterranean diet and weight loss: meta-analysis of randomised controlled trials. *Metab Syndr Relat Disord*. 2011 Feb;9(1):1-12. doi: 10.1089/met.2010.0031. Epub 2010 Oct 25. PMID: 20973675.
26. Austin G, Ferguson JJA, Garg ML. Effects of Plant-Based Diets on Weight Status in Type 2 Diabetes: A Systematic Review and Meta-Analysis of Randomised Controlled Trials. *Nutrients*. 2021 Nov 16;13(11):4099. doi: 10.3390/nu13114099. PMID: 34836356; PMCID: PMC8625212.
27. Sackner-Bernstein J, Kanter D, Kaul S. Dietary Intervention for Overweight and Obese Adults: Comparison of Low-Carbohydrate and Low-Fat Diets. A Meta-Analysis. *PLoS One*. 2015 Oct 20;10(10):e0139817. doi: 10.1371/journal.pone.0139817. PMID: 26485706; PMCID: PMC4618935.
28. Gardner CD, Trepanowski JF, Del Gobbo LC, et al. Effect of Low-Fat vs Low-Carbohydrate Diet on 12-Month Weight Loss in Overweight Adults and the Association With Genotype Pattern or Insulin Secretion: The DIETFITS Randomized Clinical Trial. *JAMA*. 2018 Feb 20;319(7):667-679. doi: 10.1001/jama.2018.0245. Erratum in: *JAMA*. 2018 Apr 3;319(13):1386. Erratum in: *JAMA*. 2018 Apr 24;319(16):1728. PMID: 29466592; PMCID: PMC5839290.
29. Bray GA, Ryan DH. Evidence-based weight loss interventions: Individualised treatment options to maximise patient outcomes. *Diabetes Obes Metab*. 2021 Feb;23 Suppl 1:50-62. doi: 10.1111/dom.14200. Epub 2020 Nov 24. PMID: 32969147.
30. Steger FL, Donnelly JE, Hull HR, Li X, Hu J, Sullivan DK. Intermittent and continuous energy restriction result in similar weight loss, weight loss maintenance, and body composition changes in a 6 month randomised pilot study. *Clin Obes*. 2021 Apr;11(2):e12430. doi: 10.1111/cob.12430. Epub 2020 Dec 10. PMID: 33305526.

LEARNING POINTS

- **To date, studies have failed to demonstrate the superiority of one diet plan over another as patients can lose weight on any diet in the short term. Long-term effectiveness studies are lacking.**
 - **Any dietary approach to weight loss should be individualised and consider the health status, personal preferences, and ability of the person to sustain the recommendations in the plan.**
 - **As healthcare professionals, we should be realistic when discussing treatment expectations with our patients and provide ongoing support to ensure long-term weight loss maintenance and manage relapse.**
-